

1. **Ebssa, L., Borgemeister, C. & H.-M. Poehling 2004a.** Effectiveness of different species/strains of entomopathogenic nematodes for control of western flower thrips (*Frankliniella occidentalis*) at various concentrations, host densities, and temperatures. *Biological Control* 29: 145-154.
2. **Ebssa, L., Borgemeister, C. & H.-M. Poehling 2004b.** Effects of post-application irrigation and substrate moisture on the efficacy of entomopathogenic nematodes against western flower thrips *Frankliniella occidentalis*. *Entomologia Experimentalis et Applicata* (in press).
3. **Ebssa, L., Borgemeister, C. & H.-M. Poehling 2004c.** Efficacy of entomopathogenic nematodes against western flower thrips *Frankliniella occidentalis* at different pupation depths. *Nematology* (in press).
4. **Ebssa, L., Borgemeister, C., Berndt, O. & H.-M. Poehling 2001a.** Efficacy of Entomopathogenic Nematodes against Soil-dwelling Life Stages of Western Flower Thrips, *Frankliniella occidentalis* (Thysanoptera: Thripidae). *Journal of Invertebrate Pathology* 78: 119-127.
5. **Ebssa, L., Borgemeister, C., Berndt, O. & H.-M. Poehling 2001b.** Impact of entomopathogenic nematodes on different soil-dwelling stages of Western Flower Thrips, *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae), in the laboratory and under semi-field conditions. *Biocontrol Science and Technology* 11: 515-525.