

### Peer reviewed publications and books

- Abu Hatab, M., Gaugler, R. & Ehlers, R.-U. 1998. Influence of culture method on *Steinernema glaseri* lipids. **Journal of Parasitology** 84, 215-221.
- Aumann, J. & Ehlers, R.-U. 2002. Physico-chemical properties and mode of action of a signal from the symbiotic bacterium *Photorhabdus luminescens* inducing dauer juvenile recovery in the entomopathogenic nematode *Heterorhabditis bacteriophora*. **Nematology** 8(3), 849-853.
- Blossey, B. & Ehlers, R.-U. 1991. Entomopathogenic nematodes (*Heterorhabditis* spp. and *Steinernema anomali*) as potential antagonists of the biological weed control agent *Hylobius transversovittatus* (Coleoptera: Curculionidae). **Journal of Invertebrate Pathology** 58, 453-454.
- Blum, B., Ehlers, R.-U., Haukeland-Salinas, S., Hokkanen, H., Jung, K., Kuhlmann, U., Menzler-Hokkanen, I., Ravensberg, W., Strasser, H., Warrior, P. & Wilson, M. 2003: Biological control agents: Safety and regulatory policy. **BioControl** 48, 477-484.
- Boemare, N., Nealon, K. & Ehlers, R.-U. 1997. *Xenorhabdus-Steinernema* and *Photorhabdus-Heterorhabditis*, entomopathogenic bacterium nematode symbioses. **Symbiosis** 22, 1-3.
- Ehlers, R.-U. 1990. Environmental and biotic factors influencing the control potential of entomopathogenic nematodes of the genus *Steinernema* and *Heterorhabditis*. **Nematologica** 36, 347.
- Ehlers, R.-U. 1996. Current and future use of nematodes in biocontrol: Practice and commercial aspects in regard to regulatory policies. **Biocontrol Science & Technology** 6, 303-316.
- Ehlers, R.-U. 1998. Entomopathogenic nematodes - Save biocontrol agents for sustainable systems. **Phytoprotection** 79, 94-102.
- Ehlers, R.-U. 2001. Mass production of entomopathogenic nematodes for plant protection. **Applied Microbiology & Biotechnology** 56, 623-633.
- Ehlers, R.-U. 2003: Biocontrol Nematodes In: H.M.T. Hokkanen & A.E. Hajek. Environmental Impacts of Microbial Insecticides – Need and Methods for Risk Assessment Kluwer Academic Publisher, Dordrecht, pp. 177-220.
- Ehlers, R.-U. 2004: Entomopathogenic nematodes in the European biocontrol market. **Communications in Agricultural and Applied Biological Sciences** 68, 3-16.
- Ehlers, R.-U., Deseö, K. V. & Stackebrandt, E. 1991. Identification of *Steinernema* spp. (Nematoda) and their symbiotic bacteria *Xenorhabdus* spp. from Italian and German soil. **Nematologica** 37, 360-366.
- Ehlers, R.-U. & Gerwien, A. 1993. Selection of entomopathogenic nematodes (Steinernematidae and Heterorhabditidae, Nematoda) for the biological control of crane fly larvae *Tipula paludosa* (Tipulidae, Diptera). **Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz** 100, 343-353.
- Ehlers, R.-U. & Hokkanen, H. M. T. 1996. Insect biocontrol with non-endemic entomopathogenic nematodes (*Steinernema* and *Heterorhabditis* spp.): Conclusions and recommendations of a combined OECD and COST workshop on scientific and regulatory policy issues. **Biocontrol Science & Technology** 6, 295-302.
- Ehlers, R.-U., Lunau, S., Krasomil-Osterfeld, K. & Osterfeld, K.-H. 1998. Liquid culture of the entomopathogenic nematode-bacterium complex *Heterorhabditis megidis/Photorhabdus luminescens*. **BioControl** 43, 77-86.
- Ehlers, R.-U. & Niemann, I. 1998. Molecular identification of *Photorhabdus luminescens* strains by amplification of specific fragments of the 16S ribosomal DNA. **Systematic & Applied Microbiology** 21, 509-519.

- Ehlers, R.-U., Niemann, I., Hollmer, S., Strauch, O., Jende, D., Shanmugasundaram, M., Mehta, U. K., Easwaramoorthy, S. K. & Burnell, A. 2000 . Mass production potential of the bacto-helminthic biocontrol complex *Heterorhabditis indica*-*Photorhabdus luminescens*. **Biocontrol Science & Technology** 10, 607-616.
- Ehlers, R.-U. & Peters, A. 1995. Entomopathogenic nematodes in biological control: feasibility, perspectives and possible risks. *In*: Hokkanen, H. M. T. & Lynch, J. M. (eds.), *Biological control: benefits and risks*. Cambridge University Press, UK, 119-136.
- Ehlers R.-U., Stoessel, S. & Wyss, U. 1990. The influence of phase variants of *Xenorhabdus* spp. and *Escherichia coli* (Enterobacteriaceae) on the propagation of entomopathogenic nematodes of the genera *Steinernema* and *Heterorhabditis*. **Revue de Nématologie** 13, 417-424.
- Ehlers, R.-U., Wulff, A. & Peters, A. 1997. Pathogenicity of axenic *Steinernema feltiae*, *Xenorhabdus bovienii* and the bacto-helminthic complex to larvae of *Tipula oleracea* (Diptera) and *Galleria mellonella* (Lepidoptera). **Journal of Invertebrate Pathology** 69, 212-217.
- Ehlers, R.-U., Wyss, U. & Stackebrandt, E. 1988. 16S RNA cataloguing and the phylogenetic position of the genus *Xenorhabdus*. **Systematic & Applied Microbiology** 10, 121-125.
- Gerritsen, L. J. M., van der Wolf, J. M., van Vuurde, J. W. L., Ehlers, R.-U., Krasomil-Osterfeld, K. C. & Smits, P. H. 1995. Polyclonal antisera to distinguish strains and form variants of *Photorhabdus* (*Xenorhabdus*) *luminescens*. **Applied & Environmental Microbiology** 61, 284-289.
- Han, R. C. & Ehlers, R.-U. 1998. Cultivation of axenic *Heterorhabditis* spp. dauer juveniles and their response to non-specific *Photorhabdus luminescens* food signals. **Nematologica** 44, 425-435.
- Han, R. & Ehlers, R.-U. 1999. Trans-specific nematocidal activity of *Photorhabdus luminescens*. **Nematology** 1, 687-693.
- Han, R. & Ehlers, R.-U. 2000. Pathogenicity, development, and reproduction of *Heterorhabditis bacteriophora* and *Steinernema carpocapsae* under axenic *in vivo* conditions. **Journal of Invertebrate Pathology** 75, 55-58.
- Han, R. & Ehlers, R.-U. 2001 . Effect of *Photorhabdus luminescens* phase variants on the *in vivo* development and reproduction of *Heterorhabditis bacteriophora* and *Steinernema carpocapsae*. **FEMS Microbiology/Ecology** 35, 239-247.
- Hazir, S., Stackebrandt, E., Lang, E., Schumann, P., Ehlers, R.-U. & Keskin, N. 2004: Two new subspecies of *Photorhabdus luminescens*, isolated from *Heterorhabditis bacteriophora* (Nematoda: Heterorhabditidae): *Photorhabdus luminescens* subsp. *kayaii* subsp. nov. and *Photorhabdus luminescens* subsp. *thracensis* subsp. nov. **Systematic and Applied Microbiology** 27, 36-42.
- Iraki, N., Salah, N., Sansour, M. A., Segal, D., Glazer, I., Johnigk, S.-A., Hussein, M. A. & Ehlers, R.-U. 2000. Isolation and characterization of two entomopathogenic nematode strains, *Heterorhabditis indica* (Nematoda, Rhabditia), from the West Bank, Palestinian Territories. **Journal of Applied Entomology** 124, 375-380.
- Jeffke, T., Jende, D., Mätje, C., Berthe-Corti, L. & Ehlers, R.-U. 2000. Growth of *Photorhabdus luminescens* in batch and fed-batch culture. **Applied Microbiology & Biotechnology** 54, 326-330.
- Jessen, P., Strauch, O., Wyss, U., Luttmann, R. & Ehlers, R.-U. 2000. Carbon dioxide triggers dauer juvenile recovery of entomopathogenic nematodes (*Heterorhabditis* spp.). **Nematology** 2, 319-324.
- Johnigk, S.-A., Ecke, F., Pöhling, M. & Ehlers, R.-U. 2004: Liquid culture mass production

- of biocontrol nematodes, *Heterorhabditis bacteriophora* (Nematoda: Rhabditida): improved timing of dauer juvenile inoculation. **Applied Microbiology and Biotechnology** 64, 651-658.
- Johnigk, S.-A. & Ehlers, R.-U. 1999a. Juvenile development and life cycle of *Heterorhabditis bacteriophora* and *H. indica*. **Nematology** 1, 251-260.
- Johnigk, S.-A. & Ehlers, R.-U. 1999b. *Endotokia matricida* in hermaphrodites of *Heterorhabditis* spp. and the effect of the food supply. **Nematology** 1, 717-726.
- Johnigk, S.-A., Hollmer, S., Strauch, O., Wyss, U. & Ehlers, R.-U. 2002. Heritability of the liquid culture mass production potential of the entomopathogenic nematode *Heterorhabditis bacteriophora*. **Biocontr. Sci. & Technol.** 12, 267-276.
- Lunau, S., Stoessel, S., Schmidt-Peisker, A. J. & Ehlers, R.-U. 1993. Establishment of monoxenic inocula for scaling up *in vitro* cultures of the entomopathogenic nematodes *Steinernema* spp. and *Heterorhabditis* spp. **Nematologica** 39, 385-399.
- Peters, A. & Ehlers, R.-U. 1994. Susceptibility of leatherjackets (*Tipula paludosa* and *T. oleracea*; Tipulidae: Nematocera) to the entomopathogenic nematode *Steinernema feltiae*. **Journal of Invertebrate Pathology** 63, 163-171.
- Peters, A. & Ehlers, R.-U. 1997. Encapsulation of the entomopathogenic nematodes *Steinernema feltiae* in *Tipula oleracea*. **Journal of Invertebrate Pathology** 69, 218-222.
- Peters, A., Gouge, D. H., Ehlers, R.-U. & Hague, N. G. M. 1997. Avoidance of encapsulation by *Heterorhabditis* spp. infecting larvae of *Tipula oleracea*. **Journal of Invertebrate Pathology** 70, 161-164.
- Premachandra, W. T. S. D., Borgmeister, C., Berndt, O., Ehlers, R.-U. & Poehling, H.-M. 2003: Combined releases of entomopathogenic nematodes and the predatory mite *Hypoaspis aculeifer* to control soil-dwelling stages of western flower thrips *Frankliniella occidentalis*. **BioControl** 48, 529-541.
- Premachandra, W. T. S. D., Borgmeister, C., Berndt, O., Ehlers, R.-U. & Poehling, H.-M. 2003: Laboratory bioassays of virulence of entomopathogenic nematodes against soil-inhabiting stages of *Frankliniella occidentalis* Pergande (Thysanoptera: Thripidae). **Nematology** 5, 539-547.
- Pütz, J., Meinert, F., Wyss, U., Ehlers, R.-U. & Stackebrandt, E. 1990. Development and application of oligonucleotide probes for molecular identification of *Xenorhabdus* species. **Applied & Environmental Microbiology** 56, 181-186.
- Rainey, F. A., Ehlers, R.-U. & Stackebrandt, E. 1995. Inability of the polyphasic approach to systematics to determine the relatedness of the genera *Xenorhabdus* and *Photorhabdus*. **International Journal of Systematic Bacteriology** 45, 379-381.
- Schroer, S. & Ehlers, R.-U. 2004. Foliar application of the entomopathogenic nematode *Steinernema carpocapsae* for biological control of Diamondback Moth larvae (*Plutella xylostella*). **Biological Control**, in press
- Schroer, S., Yi, X. & Ehlers, R.-U. 2004. Evaluation of adjuvants for foliar application of *Steinernema carpocapsae* against larvae of the diamondback moth (*Plutella xylostella*). **Nematology**, in press
- Schroer, S., Ziermann, D. & Ehlers, R.-U. 2004. Mode of action of a surfactant-Polymer-Formulation to support performance of the entomopathogenic nematode *Steinernema Carpocapsae* for control of diamondback moth larvae (*Plutella xylostella*). **Biocontrol Science and Technology**, in press
- Simoës, N., Boemare, N. & Ehlers, R.-U. 1998. COST 819 - Entomopathogenic nematodes: Pathogenicity of entomopathogenic nematodes versus insect defence mechanisms: Impact on selection of virulent strains. European Commission, Luxembourg.
- Stackebrandt, E., Ehlers, R.-U. & Rainey, F. A. 1997. *Xenorhabdus* and *Photorhabdus*: Are

- they sister genera or are their members phylogenetically intertwined. **Symbiosis** 22, 59-65.
- Strauch, O. & Ehlers, R.-U. 1998. Food signal production of *Photorhabdus luminescens* inducing the recovery of entomopathogenic nematodes *Heterorhabditis* spp. in liquid culture. **Applied Microbiology & Biotechnology** 50, 369-374.
- Strauch, O. & Ehlers, R.-U. 2000. Influence of the aeration rate on the yields of the biocontrol nematodes *Heterorhabditis megidis* in monoxenic liquid cultures. **Applied Microbiology & Biotechnology** 54, 9-13.
- Strauch, O., Niemann, I., Neumann, A., Schmidt, A. J., Peters, A. & Ehlers, R.-U. 2000. Storage and formulation of the entomopathogenic nematodes *Heterorhabditis indica* and *H. bacteriophora*. **BioControl** 45, 483-500.
- Strauch, O., Oestergaard, J., Hollmer, S. & Ehlers, R.-U. 2004. Genetic improvement of the desiccation tolerance of the entomopathogenic nematode *Heterorhabditis bacteriophora* through selective breeding. **Biological Control** 31, 218-226.
- Strauch, O., Stoessel, S. & Ehlers, R.-U. 1994. Culture conditions define automictic or amphimictic reproduction in entomopathogenic rhabditid nematodes of the genus *Heterorhabditis*. **Fundamental & Applied Nematology** 17, 575-582.
- Sulistyanto, D. & Ehlers, R.-U. 1996. Efficacy of the entomopathogenic nematodes *Heterorhabditis megidis* and *Heterorhabditis bacteriophora* for the control of grubs (*Phyllopertha horticola* and *Aphodius contaminatus*) in golf course turf. **Biocontrol Science & Technology** 6, 247-250.
- Susurluk, A., Dix, I., Stackebrandt, E., Strauch, O. & Ehlers, R.-U. 2002. Identification and ecological characterization of three entomopathogenic nematode-bacterium complexes. **Nematology** 3, 833-841.
- Tallosi, B., Peters, A. & Ehlers, R.-U. 1995. *Steinernema bicornutum* sp.n. (Rhabditida: Nematoda) from Vojvodina, Yugoslavia. **Russian Journal of Nematology** 3, 71-79.