

## **PUBLICATIONS**

- Van Tol, R.W.H.M. and M.J. Raupp (2004) Nursery and tree application (Chapter 9). In: *Nematodes as biological control agents* (Eds. P.S. Grewal, R-U. Ehlers and D.I. Shapiro-Ilan). CABI Publishing. In press.
- Van Tol, R.W.H.M., N. van Dijk and M.W. Sabelis (2004). Host plant preference and performance of the vine weevil *Otiorhynchus sulcatus*. *Agricultural and Forest Entomology*. In press.
- Van Tol, R.W.H.M. and R.L. Gwynn (2004). Field Efficacy (Chapter 5). In: Quality control of entomopathogenic nematodes. *Proceedings of COST 819 Workshop "Application and Formulation of Entomopathogenic Nematodes"*, Eds. J.M. Grunder. In press.
- Van Tol, R.W.H.M., J.H. Visser and M.W. Sabelis (2004). Behavioural responses of the vine weevil, *Otiorhynchus sulcatus*, to semiochemicals from conspecifics, *Otiorhynchus salicicola*, and host plants. *Entomologia Experimentalis et Applicata* **110**, 145-150.
- Van Tol, R.W.H.M. (2002). Fatal attraction – Novel strategies for vine weevil control. *PhD thesis*, University of Amsterdam, the Netherlands, ISBN 90 76894 21 3
- Van Tol, R.W.H.M., J.H. Visser and M.W. Sabelis (2002). Olfactory responses of the vine weevil, *Otiorhynchus sulcatus*, to tree odours. *Physiological Entomology* **27**, 213-222.
- Van Tol, R.W.H.M. and J.H. Visser (2002). Olfactory antennal responses of the black vine weevil (*Otiorhynchus sulcatus*) to plant volatiles. *Entomologia Experimentalis et Applicata* **102**, 49-64.
- Van Tol, R.W.H.M., A.T.C. van der Sommen, J. van Bezooijen and P.H. Smits (2001). Olfactory attraction of *Heterorhabditis megidis* to roots and vine weevil feeding-induced root semiochemicals. In: *Developments in entomopathogenic nematode/bacterial research* (eds. C. T. Griffin, A. M. Burnell, M. J. Downes and R. Mulder). *COST Action 819, Proceedings of the workshop at the National University of Ireland Maynooth, 13-15 April 2000, EUR 19696*, 156-163.
- Boff, M.I.C., R.W.H.M. Van Tol and P.H. Smits (2001). Behavioural response of *Heterorhabditis megidis* towards plant roots and insect larvae. *Biocontrol* **47**, 67-83.
- Van Tol, R.W.H.M., A.T.C. van der Sommen, M.I.C. Boff, J. van Bezooijen, M.W. Sabelis and P.H. Smits (2001). Plants protect their roots by alerting the enemies of grubs. *Ecology Letters* **4**, 1-3.
- Van Tol, R.W.H.M., J.H. Visser and M.W. Sabelis (2000). Responses of the black vine weevil (*Otiorhynchus sulcatus*) to weevil and host-plant odours. *Proc. Exper. & Appl. Entomology, N.E.V. Amsterdam*, **11**, 109-114.
- Van Tol, R.W.H.M. and A.C. Schepman (1999). Influence of host and plant roots on the migration of *Heterorhabditis* sp. (NWE) in peat soil. *Proceedings of COST 819 Workshop "Application and Persistence of entomopathogenic nematodes"*, Todi, Italy, 16-20 May 1995, 117-121.

- Van Tol, R.W.H.M. and J.H. Visser (1998). Host plant preference and antennal response of the black vine weevil (*Otiorhynchus sulcatus*) to plant volatiles. *Proc. Exper. & Appl. Entomology, N.E.V. Amsterdam*, **9**, 35-40.
- Van Tol, R.W.H.M., Van Bezooijen, J. and T.A.C.M. Ketelaars (1998). Searching behaviour of entomopathogenic nematodes: Roots and soil temperature determine success of black vine weevil (*Otiorhynchus sulcatus*) control. *IOBC/WPRS Bulletin Insect Pathogens and Insect Parasitic Nematodes* **21** (4), 187-191.
- Van Tol, R.W.H.M. (1997). Biological and chemical control of the vine weevil *Otiorhynchus sulcatus*. *BPO report* **50**, 46 p.
- Dolmans, N.G.M. and R.W.H.M. Van Tol (1996). Prospects for chemical control of black vine weevil (*Otiorhynchus sulcatus*) in nursery stock. *Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft* **316**, 108-112.
- Van Tol, R.W.H.M. (1996). A strategy for control of black vine weevil (*Otiorhynchus sulcatus*) in an Integrated Pest Management programme in nursery stock. *Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft* **316**, 76-80.
- Van Tol, R.W.H.M. (1996). Prospects for biological control of black vine weevil (*Otiorhynchus sulcatus*) in nursery stock. *Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft* **316**, 69-75.
- Van der Horst, M.J. and R.W.H.M. Van Tol (1995). Integrated Pest Management in nursery stock in The Netherlands. *Med. Fac. Landbouww. Univ. Gent* **60/3a**, 759-762.
- Van Tol, R.W.H.M. (1994). Influence of temperature on the control of the black vine weevil with strains of some insect-parasitic nematodes. *IOBC/WPRS Bulletin* **17(3)**, 116-119.
- Van Tol, R.W.H.M. (1993). Efficacy of control of the black vine weevil (*Otiorhynchus sulcatus*) with strains of *Heterorhabditis* sp., *Steinernema* sp. and the fungus *Metarhizium anisopliae* in nursery stock. *Med. Fac. Landbouww. Univ. Gent* **58/2a**, 461-467.
- Van Tol, R.W.H.M. (1993). Control of the black vine weevil (*Otiorhynchus sulcatus*) with different isolates of *Heterorhabditis* sp. and *Metarhizium anisopliae* in nursery stock. *Proc. Exper. & Appl. Entomolgy, N.E.V. Amsterdam*, **4**, 181-186.